

**RESEARCH ARTICLE****Vitamin Contents in Five Edible Varieties of Mushroom Consumed in Umuagwo Community, Imo State, Nigeria**Ibe CC¹, CN Osuji¹, TO Akunna¹, EO Ahaotu² and EU Nwabueze¹¹Department of Science Laboratory Technology, Imo State Polytechnic Umuagwo, Nigeria²Department of Animal Production Technology, Imo State Polytechnic Umuagwo, Nigeria**ARTICLE INFO**

Received: April 23, 2013
Revised: June 23, 2013
Accepted: August 16, 2013

Key words:

Mushrooms
Nutrient supply
Umuagwo
Vitamin composition

***Corresponding Address:**

Ibe CC
vieng663@hotmail.com
emmaocy@yahoo.com

ABSTRACT

The vitamin contents of five wild growing mushroom species in Umuagwo Community, Imo State, Nigeria were investigated. The aim of this study was to assess and compare the nutritional contents of various species of mushrooms found in Umuagwo Community. Of all the five wild mushroom species studied, vitamins A, C, E and niacin were found to be relatively higher in *Lactarius deliciosus*. This was followed by *Daedalea elegans* (P<0.05). Thiamin was also, relatively higher in *Daedalea elegans*, followed by *Lactarius deliciosus*, *Coprinus africanus*, *Pleurotus florida* and *Lepiota procera* (P<0.05). Vitamins A and E were only found in *Agaricus bisporus*. Based on these results, the mushroom species are good sources of vitamin C, vitamin B1 and niacin and could be considered ideal supplements for many low vitamin food materials in our diets.

Cite This Article as: Ibe CC, CN Osuji, TO Akunna and EO Ahaotu, 2013. Vitamin contents in five edible varieties of mushroom consumed in Umuagwo Community, Imo State, Nigeria. *Inter J Agri Biosci*, 2(5): 177-180. www.ijagbio.com

INTRODUCTION

Mushroom is a macro fungus with a distinctive fruiting body, which can be either epigeous or hypogeous, and large enough to be seen with naked eye and to be picked by hand (Chang and Miles, 1992). They lack chlorophyll present in plants. Mushrooms represent one of the world's greatest untapped resources of nutritious food. Mushrooms are classified under sub division *Basidiomycotina*. They grow on dead and decaying organic material. From these decaying substrates, they absorb their nutrients with the help of very fine thread like structures (mycelium) which penetrates into substratum and generally not visible on the surface. After the mycelium has grown profusely and absorbs sufficient food materials, it forms the reproductive structures which generally comes out of the substrate and forms fruiting body, commonly known as mushrooms. The mushrooms fruiting body may be umbrella like or of various shapes, size and colours. All mushrooms are not edible and some are highly poisonous species which are commonly referred to as 'toad stools' (Crisian and Sands, 1978). Of several thousand mushroom species known worldwide, only around 2000 are considered edible, of which about 20 are cultivated commercially with only 4 to 5 under industrial production (Chang, 1990). Cultivation of the

saprophytic edible mushrooms may be the only and currently economical biotechnology for lignocelluloses organic waste recycling that combines the production of protein rich food with the reduction of environmental pollution (Obodai *et al.*, 2003). Mushrooms are rich in protein, minerals, vitamins, fibers and they contain an abundance of essential amino acids (Sadler, 2003; Chadha and Sharma, 1995). Therefore, mushrooms can be a good supplement to cereals (Chang and Buswell, 1996). Mushroom is being widely used as food and food supplements from ancient times. They are increasingly being recognized as one of the important food items for their significant roles in human health, nutrition and diseases (Chang, 1996). Mushrooms are recognized as the alternative source of good quality protein and are capable of producing the highest quantity of protein per unit area and time from the worthless agro-wastes (Chadha and Sharma, 1995). Mushrooms can substantiate the sufferings from malnutrition to some extent, because they produce large quantities in a short time and provide more protein per unit area than other crops (Hossain *et al.*, 2007). They are low in calories, carbohydrates, calcium and sodium. They contain high proportion of unsaturated fat but virtually no harmful lipid or cholesterol. It contains a large amount of vitamins such as Thiamine 1.4-2.2 mg (%), Riboflavin 6.7-9.0 mg (%), Niacin 60.6-73.3 mg (%),

Biotin, Ascorbic acid 92-144 mg (%), Pantothenic acid 21.1-33.3 mg (%) and Folic acid 1.2-1.4 mg/100g in dry weight basis (Hossain *et al.*, 2007). The minerals found in mushroom are calcium, Iron, Manganese, Magnesium, Zinc, Selenium etc (Alam *et al.*, 2007). The mineral proportions vary according to the species, age and the diameter of the fruiting body. It also depends upon the type of the substratum (Demirbas, 2001). The mineral content of wild edible mushrooms has been found higher than cultivated ones (Aletor, 1995; Mattilla *et al.*, 2001; Rudawska and Leski, 2005). Mushroom has 1000 of variety to enrich the food basket of human health. It is an alternative rich source of meat, fish, vegetables, fruits etc. Mushroom's nutritional value proves to have many health benefits.

Consequently, upon the nutritional value of mushrooms, they have been also recognized as medicines for effective treating of cancer, cholesterol reduction, stress, insomnia, asthma, allergies and diabetes (Bahl, 1983). Also, medicinal properties, such as antifungal, antibacterial, antioxidant, antiviral anti-hypotensive, renal protective, immune modulatory and anticancer activities of mushroom extracts have been reported (Chang and Miles, 2004; Musieba *et al.*, 2011). Bahl (1983) reported that mushrooms cure epilepsy, wounds, skin diseases, heart ailments, rheumatoid arthritis, Cholera, fevers, diaphoretic, diarrhea, dysentery, cold, anesthesia, liver disease, gall bladder diseases and can be used as vermicides. The main therapeutically active components were found to be polysaccharides, especially β -D- glucans (Wani *et al.*, 2010 and Musieba *et al.*, 2011). Most of these medicinal components are now available in tablet and capsule forms, which are called mushroom nutraceuticals and are used as food supplements (Yang *et al.*, 1993). Extracts from certain mushroom species (*Ganoderma lucidum*, *Ganoderma lucidium* and *Lentinus tigrinus*) have been used to lower blood pressure and serum cholesterol in hypertensive rats (Kabir *et al.*, 1988; Ren *et al.*, 1989). Thousands of years ago, fructifications of higher fungi have been used as a source of food (Mattila *et al.*, 2001) due to their chemical composition which is attractive from the nutrition point of view. During the early days of civilization, mushrooms were consumed mainly for their palatability and unique flavors (Rai, 1994). Present use of mushrooms is totally different from traditional because, lot of research has been done on the chemical composition of mushrooms, which revealed that mushrooms can be used as a diet to combat diseases. The early history regarding the use of mushrooms in different countries has been reviewed by number of workers (Oyetayo *et al.*, 2011; Bilal *et al.*, 2010; Tibuhwa *et al.*, 2011). Ayodele *et al.* (2011) recommended that 100 to 200 g of mushrooms (dry weight) is required to maintain an optimal nutritional balance in a man weighing 70 kg. Bano (1976) suggested that food value of mushrooms lies between meat and vegetables. Crisan and Sands (1978) observed that mushrooms generally contain 90% water and 10% dry matter. More so, the protein content varies between 27 and 48%. Carbohydrates are less than 60% and lipids are between 2 to 8%. Gruen and Wong (1982) indicated that edible mushrooms were highly nutritional and compared favourably with meat, egg and milk. This study was aimed at characterizing

some wild growing edible mushrooms common in Umuagwo Community, Eastern Nigeria for vitamin compositions as an assessment of their nutritional values. The study was also aimed at providing useful information for selecting mushroom species that are rich in these nutrients for artificial production, since rapid deforestation is almost sending these important food resources into extinction in the region.

MATERIALS AND METHODS

Determination of vitamin composition

The five mushroom species used in this study were harvested from Umuagwo Southeast Nigeria. The mushrooms, after collection, were washed cleaned with distilled water, cut into small pieces and ground using electric blender to homogenize the sample. The vitamin contents of the edible mushroom varieties from Umuagwo, Imo State were determined according to the method described by Onwuka (2005).

RESULTS AND DISCUSSION

Vitamin analysis was carried out on five wild grown edible mushroom varieties consumed in Umuagwo, South Eastern Nigeria; *Daedalea elegans*, Bell like mushroom, *Lactarius deliciosus*, Hard tissue mushroom, *Lepiota procera*, Meat like mushroom, *Pleurotus florida*, Tree like mushroom and *Coprinus africanus*, Palm mushroom. Results of the vitamin compositions are presented in Table 1. The vitamin contents of all the mushroom varieties varied significantly among the species ($P < 0.0001$). Vitamin A contents of the mushrooms varied from 0-38.36% with a mean value of $38.36 \pm 0.18\%$. Vitamin A was only detected in *Lactarius deliciosus* at a mean concentration of 38.36mg/100g while there was no vitamin A in other mushroom varieties studied. The value obtained for vitamin A in *Lactarius deliciosus* was significantly higher than most animal and plant materials known to contain significant amount of vitamin A including veal liver (28mg/100g), butter (0.59mg/100g), cheese (0.39mg/100g), egg (0.28), salmon (0.041mg/100g) and milk (0.04mg/100g) (Souci *et al.*, 2000). Vitamin A is a generic term for a group of lipid soluble compounds related to retinol. The main function of vitamin A has been found to include proper vision, reproduction, growth and development, cellular differentiation and immune function. Its earliest deficiency symptom is impaired dark adaptation or night blindness. Severe deficiency could cause Xerophthalmia, characterized by changes in the cells of the cornea that ultimately result in corneal ulcers, scarring and blindness (Bowman, 2001).

The value obtained for vitamin C (ascorbic acid) indicated that the mushrooms are considerable sources of vitamin C. The amount of vitamin C spanned from 0.68-1.70% with a mean concentration of $7.67 \pm 15.88\%$. The highest vitamin C content was found in *Lactarius deliciosus* while the lowest concentration of vitamin C was found in coral mushroom. Sapers *et al.*, 1999 and Mattila *et al.*, 2001 reported vitamin C content of up to 7mg/100g of edible parts of mushrooms. The values of vitamin C obtained for each of the mushroom varieties

Table 1: Vitamin contents of five edible mushroom varieties in Umuagwo Community, Imo State, Nigeria.

Sample	Vit. A (mg/100g)	Vit.C (mg/100g)	Vit.E (IU/100g)	Thiamine (mg/100g)	Niacin (mg/100g)
<i>Daedalea elegans</i>	0.00±0.00 ^b	1.04±0.01 ^b	0.00±0.00 ^b	0.52±0.57 ^a	1.31±0.04 ^b
<i>Coprinus africanus</i>	0.00±0.00 ^b	0.74±0.01 ^c	0.00±0.00 ^b	0.19±0.02 ^c	0.66±0.01 ^d
<i>Lactarius deliciosus</i>	38.36±0.18 ^c	1.70±0.01 ^a	5.70±0.65 ^a	0.26±0.01 ^b	1.65±0.00 ^a
<i>Pleurotus florida</i>	0.00±0.00 ^b	0.70±0.01 ^d	0.00±0.00 ^b	0.07±0.01 ^d	0.33±0.01 ^c
<i>Lepiota procera</i>	0.00±0.00 ^b	0.68±0.58 ^d	0.00±0.00 ^b	0.27±0.02 ^b	0.98±0.01 ^c
Grand Average	7.67±15.88	0.97±0.46	1.14±2.37	0.26±0.27	0.99±0.48
P. value	P<0.0001	P<0.0030	P<0.0001	P<0.3445	P<0.0001
n- value	n=3	n=3	n=3	n=3	n=3

studied were significantly lower than the reported value. The value of vitamin C content obtained in *Lactarius deliciosus* (1.70mg/100g) compares favourably with value of 6.9mg/100g reported respectively on *Lactarius deliciosus* and *Coprinus africanus* by Mattila *et al* (2001) and Mattila *et al* (2002). Okhuoya *et al* (2010) reported a significantly higher vitamin C content of 8.6mg/100g in *Lactarius deliciosus*. The value of vitamin C obtained in *Coprinus africanus* (0.74mg/100g) was significantly lower than the vitamin C content of 1.6mg/100g reported on *Coprinus africanus* by Mattila *et al* (2001) and Mattila *et al* (2002). The result obtained for vitamin E indicates that the mushroom varieties studied are significantly low in vitamin E. Only *Lactarius deliciosus* recorded a detectable amount of vitamin E content of 5.70 IU/100g among the studies mushroom varieties. There dearth of information on the vitamin E content of edible wild mushroom varieties. However, the values obtained are in concordant with the reports on available literature that mushrooms are significantly low in vitamin E (Outila *et al.*, 1999). Studies have shown that vitamin E decrease symptoms of premenstrual syndrome and certain types of breast disease (Comb, 2001), serves as a useful supplement for some neurological diseases such as Alzheimer's disease (Reboul *et al.*, 2006). Serves as a major fat soluble antioxidant of the body and also carries out Non-antioxidant functions in cell signaling, gene expression and regulation of other cell functions (FSA, 2003). Thiamine is a water-soluble B-complex vitamin. It is required in significant amount for the vital functions it perform such as the flow of electrolytes in and out of nerve and muscle cells (through ion channels), coenzyme for both Co-enzyme in energy metabolism and for pentose metabolism (as a basis for nucleic acids), nerve impulse conduction and muscle action; as part of treatment for metabolic disorders like maple syrup urine disease, for treatment of beri beri among others (Wakita, 1976). The values 0.19-0.54% obtained for the mushroom varieties that the mushrooms are low in thiamin compared to yeast (Wakita, 1976). The opinion is that, with respect to thiamin content, mushrooms are a bridge between yeast and other food products of vegetable origin (Karchocha and Mlodeoki, 1965). The low thiamin content in mushrooms when compared to yeast could be accounted for by the presence of thiaminase, an enzyme that destroy thiamin (Wakita, 1976). Apparently, mushrooms have the hydrolase and a few also have the transferase, both of which destroy the vitamin (Kurtzman, 2005). However, the thiaminase can be destroyed during normal processing and cooking procedures. It is possible that with proper care, the thiamin in the mushrooms could be greatly increased (Kurtzman, 2005).

The presence of niacin was observed in significantly varying amounts in the five edible mushroom varieties

that were studied. The result of the analysis showed that niacin content of the mushrooms varied from 0.3-1.65% with a mean value of 0.99±0.48. According to Kurtzman, 2005, niacin is very high in mushrooms. It is the anti-pellagra vitamin, but it is also recommended for controlling blood cholesterol (Kurtzman, 2005). It provides 23-26% of the recommended daily intake (sRDI) for niacin per 100g (Surabhi and Devina, 2011). Watt and Merrill (1963), Okhuoya *et al.* (2010), Mattilla *et al.* (2001; 2002) reported a significantly higher concentration of niacin in *Lactarius deliciosus*. They also reported a significantly higher amount of niacin in *Coprinus africanus*. However, the results obtained for the studied mushroom varieties suggest a positive correlation between niacin and vitamin A ($r = 0.71$), niacin and vitamin C ($r = 0.74$), niacin and vitamin E ($r = 0.7$), and niacin and thiamin ($r = 0.38$).

Conclusion

The study indicated that some of the mushrooms, particularly *Lactarius deliciosus* are good sources of the vitamins analyzed and could be ideal supplements for many low vitamin containing food materials including meat in our densely populated country.

REFERENCES

- Aletor VA, 1995. Compositional studies on edible tropical species of mushrooms. *Food Chemistry* 54: 265-268.
- Bahl N, 1983. Medicinal value of edible fungi. *In: Proceeding of the International Conference on Science and Cultivation Technology of edible Fungi. Indian Mushroom Science, II: 203-209.*
- Bano Z, 1976. Nutritive value of Indian mushrooms and medicinal practices. *Eco Bot*, 31: 367-371.
- Bowman BA and RB Russell, 2001. Present Knowledge in Nutrition. International Life Science Institute. Press Washington, DC 8pp.
- Oyetayo OV, 2011. Medicinal Uses of Mushrooms In Nigeria: Towards Full And Sustainable Exploitation. *Afrin J Tradit, Complementary Alternative Med*, 8: 267-274.
- Chadha KL and SR Sharma, 1995. Mushroom research in India-History, Infrastructure and Chadha KL and Sharma SR (eds) Malhotra publishing house, New Delhi, 537-551.
- Chang ST, 1996. Mushroom research and development-equality and mutual benefit. In Chang ST and Buswell JA. *Mushroom nutraceuticals. World J Microbiol Biotechnol*, 12: 473-476.
- Chang ST and PG Miles, 1992. Mushroom biology-a new discipline. *The Mycologist*, 6:64-65.
- Chang ST and PG Miles, 2004. Mushrooms: Cultivation, Nutritional Value, Medicinal Chang ST. Future trends

- in cultivation of alternative mushrooms. *Mushroom J*, 2004; 215: 422-423.
- Musieba F, Okoth S and RK Mibey, 2011. First Record of Occurrence of *Pleurotus citrinopileatus* Singer on new hosts in Kenya. *Agric Biol J North Am*, 2: 1304-1309.
- Crisan EV and Sands A Nutritional value 1978, 2001. *In: The Biology and Cultivation of Edible Demirbas A. Concentrations of 21 metals in 18 species of mushrooms growing in the east Black Sea region. Food Chemistry* 198, 2001, 75: 453-457.
- Okhuoya JA, EO Akpaja, OO Osemwegie, AO Oghenekaro and CA Ihayere, 2010. Nigeria Mushrooms: Underutilized Non-Wood Forest Resources. *J Appl Environ Manage*, 14: 43-54.
- Eurabhi S and V Devina, 2011. White button mushroom (*Agaricus bisporus*): Composition, nutritive value, shelf-life extension and value addition. *Inter J Food Fermentation Technology* 1: 185-199.
- Gruen, VE and HX Wong, 2003. Immunomodulatory and Antitumour activities of a polysaccharide-peptide complex from a mycelial culture of *Trichoderma* sp. *Sciences*, 57: 269-281.
- Hossain MS, Alam N, Amin SMR, Basunia MA and Rahman A.2007. Essential Fatty Acid Contents of *Pleurotus ostreatus*, *Ganoderma lucidum* and *Agaricus bisporus*. *Bang J Mushroom*, 1: 1-7.
- Osemwegie OO, JA Okhuoya, AO Oghenekaro and GA Evueh, 2010. Macrofungi Community in Rubber Plantation and a Forest of Edo State, Nigeria. *J Appl Sci*, 10: 391-398.
- Kabir Y, S Kimura and T Tamura, 1988. Dietary effect of *Ganoderma lucidum* mushroom on blood pressure and lipid levels in spontaneously hypertensive rats (SHR). *J Nutr Sci Vitaminol*, 34: 433-438.
- Karchocha I and H Mlodeoki, 1965. *Tropical macrofungi*. Macmillan Press, London, 110pp.
- Kurtzman RH, 2005. Mushrooms: sources for modern Western medicine. *Micologigia Applicada Internation* 17: 21-33.
- Ayodele SM, FP Emmaneul and OF Agianaku, 2011. Comparative Studies on the Effect of sun, smoke, and oven drying methods on the nutrient contents of four wild edible mushrooms in Nigeria. *J Natural Prod Plant Res*, 1: 70-74.
- Mattila P, K Konko, M Euroala, JM Pihlava, J Astola, L Vahteristo, V Hietaniemi, J Kumpulainen, M Valtonen and V Piironen, 2001. Contents of vitamins, mineral elements and some phenolic compounds in cultivated mushrooms. *J Agri Food Chem*, 49: 2343-2348.
- Obodai M, J Cleland-Okine and KA Vowotor, 2003. Comparative study on the growth and yield of *Pleurotus ostreatus* mushroom on different lignocellulosic by-products. *J Indust Microbiol Biotechnol*, 30: 146-149.
- Outila T, P Marttila, V Piironen and CJ Lamberg-Allardt, 1999. Bioavailability of vitamin D from wild edible mushrooms (*cantharellus tubaeformis*) as measured with a human bioassay. *Amer J Clinic Nutr*, 69: 95-98.
- Rai RD, 1994. Nutritional and medicinal values of mushrooms. *In: Advances in Horticulture*.
- Ren L, AV Visitev, AN Grekhov, VV Tertov and VA Tutelyan, 1989. Antiatherosclerotic properties of macrofungi. *Voprosy Pictaniya* 189; 1: 16-19.
- Bilal AW, RH Bodha and A Wani, 2010. Nutritional and Medicinal Importance of Mushrooms. *J Med Plants Res*, 4: 2598-2604.
- Rudawska M and T Leski, 2005. Macro and micro elemental contents in fruiting bodies of wild mushrooms from the Netecka forest in west-central Poland. *Food Chem*, 92: 499-502
- Sadler M, 2003. Nutritional properties of edible fungi. *British Nutritional Foundation and Nutritional Bulletin*, 28: 305-308.
- Sapers GM, RL Miller, SW Choi and PH Cooke, 1999. Structure and composition of mushrooms as affected by hydrogen peroxide wash. *J Food Sci*, 64: 889-892.
- Tibuhwa DD, MN Muchane, CW Masiga, C Mugoya and M Muchai, 2011. An Inventory of Macro-Fungi and their Diversity in the Serengeti-Masai Mara Ecosystem, Tanzania and Kenya. *J Biol Sci*, 11: 399-410.
- Wakita S, 1976. Thiamine-destruction by mushrooms. *Science Report of the Yokohoma National University Section 2*: 39-70.
- Wani BA, RH Bodha and AH Wani, 2010. Nutritional and medicinal importance of mushrooms. *J Med Plants Res*, 4: 2598-2604.
- Watt BK and AL Merrill, 1963. *Composition of Foods. Agricultural Handbook No. 8. US. Department of Agriculture, Washington, DC, 190pp.*
- Yang QY, YJ Hu, XY Li, SX Yang, JX Liu, TF Liu, GM Xu and LM Liao, 1993. A New Biological Response Modifier - PSP. *In: Chang ST, Buswell JA and Chiu, SW (eds.) Mushroom Biology and Mushroom Products. The Chinese University Press, Hong Kong, 247-259pp.*