

RESEARCH ARTICLE**Effect of Different Concentrations of Metal Ions on *Bacillus* and *Pseudomonas* spp. Isolated from Industrial Effluents of Aluminium Extrusion Industry (ALEX), Inyishi, Imo State, Nigeria**JN Amaechi¹, JA Nwodu¹ and EO Ahaotu²¹Department of Science Laboratory Technology, Imo State Polytechnic Umuagwo, Nigeria²Department of Animal Science and Health Technology, Imo State Polytechnic Umuagwo, Nigeria**ARTICLE INFO**

Received: January 04, 2014
Revised: February 15, 2014
Accepted: March 18, 2014

Key words:

Bacillus metal effect
Optical density
Pseudomona

***Corresponding Address:**

JN Amaechi
vieng663@hotmail.com

ABSTRACT

Effects of different metals on growth of bacteria were identified to be heavy metal resistant. A total of six bacterial samples were selected, re-purified and re-identified from the bacterial isolates. Different morphological and biochemical tests were performed that differentiated bacteria into two genera (*Bacillus* spp. and *Pseudomonas* spp). Bacteria were grown at pH 7 and temp of 37°C up to incubation time of 96 hours. Three metals (Na⁺, Ca²⁺, Zn²⁺) were used with concentrations ranged between 25-200 mg/ml. It was found that as the concentrations of metals increased, growth of *Bacillus* and *Pseudomonas* spp. firstly increased and then decreased in case of all selected metals. Lowest growth occurred at 200 mg/ml of metal concentration. Optical density at 600 nm was considered as a measure of growth of bacteria. As growth increased, optical density increased and vice versa. Na⁺ was found to have a more negative effect on growth of bacterial isolates as compared to other two metals that is, Ca²⁺, Zn²⁺. *Bacillus* spp. Under investigation was found to be more resistant to these metals as compared to *Pseudomonas* spp. Both bacterial isolates showed their maximum growth after 72 hours of incubation in most of the cases, afterwards growth decreased. Decreasing resistance patterns for both of bacterial species were as follows; Ca²⁺ > Zn²⁺ > Na⁺. The findings revealed the potential application of *Bacillus* and *Pseudomonas* spp. for the removal of heavy metals from industrial wastewaters under the conditions where salts of Sodium, Zinc and Calcium are present.

Cite This Article as: Amaechi JN, JA Nwodu and EO Ahaotu, 2014. Effect of different concentrations of metal ions on *bacillus* and *Pseudomonas* spp. isolated from industrial effluents of aluminium extrusion industry (ALEX), Inyishi, Imo State, Nigeria. Inter J Agri Biosci, 3(2): 55-60. www.ijagbio.com

INTRODUCTION

Increase in global population lead to variation in natural and anthropogenic activities which encourages the contamination of various terrestrial and aquatic ecosystems with heavy metals, inorganic, organic compounds and radio nuclides (Rani, 2003). The effluents discharged by different industries contain higher values of physio-chemical parameters like temperature, pH, conductivity, hardness, alkalinity, chemical oxygen demand, total soluble salts, Nitrites and cations (Na, K, Ca and Mg) (Latif *et al.*, 2008). Discharge of metals and some non-metals into water bodies have serious environmental effects. In Nigeria, the main contributors to the surface and ground water pollution are the by-products of various industries such as textile, metal, dying chemicals, fertilizers, pesticides, cement, petrochemical,

energy and power, leather, sugar processing, construction, steel, engineering, food processing, mining and others (Tariq *et al.*, 2006). Main reason for this metal contamination is that 78% of the land is irrigated by wastewater generated from industry. The Inyishi River, Imo State, Nigeria and its tributaries provide water to over 7.1 million hectares of land in the country. A rapidly growing population, saline groundwater, a poorly performing irrigation distribution system, and recurrent droughts have led to increased water shortages. Under these conditions, the use of untreated urban wastewater for agriculture has become a common and widespread practice (Ensink *et al.*, 2002). Heavy metals are major pollutants in marine, ground, industrial and even treated wastewater (Valdman *et al.*, 2001). The alarming situation has aroused in the urban areas due to careless disposal of industrial waste and faulty drainage system that has badly

affected the quality of the river water (Ugochukwu, 2004; Chindah *et al.*, 2004; Emongor *et al.*, 2005). Major contributors to heavy metal water pollution are tanneries, brewery, textile, pottery electroplating, metal finishing, mining, dyeing and printing industries, ceramic, photographic and pharmaceutical industries, scattered in the region (Azzaoui *et al.*, 2002; Vutukuru, 2003). Bioremediation, which is essentially the use of microbial metabolism, seems to offer a viable, safer, more efficient and less expensive alternative to physiochemical methods for pollution treatment (Pandey and Jain, 2002). Microorganisms are of primary importance in bioremediation of soils and waters contaminated with heavy metals, essentially because of their ability to alter the chemical status of the metal ions and in turn the metal ion mobility through processes such as reduction, accumulation, mobilization and immobilization. In the treatment of waste water microorganisms (mainly bacteria) including both the gram-positive types, such as *Bacillus* spp. and the gram negative types such as *Pseudomonas* sp. use the soluble organic compound and convert them into carbon dioxide, water and energy to produce new cell (Minna *et al.*, 2006). Interestingly, microorganisms do require certain metal ions, like those of Cu^{2+} , Zn^{2+} , Ni^{2+} , in very low concentration as essential micronutrients as components of important cofactors in enzymatic reactions. However at higher concentrations, researchers have reported that these metals may cause cellular death by intervening with nucleic acids and enzyme activation sites (Sani *et al.*, 2001). High levels of TDO (Total dissolved oxygen) and Sodium Chloride is typically present in tannery effluent.

The TDO concentration can reach 7,000 mg/l and in some cases more than 15,000 mg/l. Effluent of brewery industry, sugar industry, effluent of the fluoride removal process of the electronics industry, fleshing wastes, textile industry, oil industry, marble industry, metal finishing company contain high concentration of calcium (Barefoot *et al.*, 2002). Zinc is an important trace metal used in numerous biological processes, however, at higher concentration, Zinc becomes a biological hazard. Therefore there is significant interest regarding Zinc removal from wastewater and its toxicity for humans at levels of 100–500 mg/day. Addition of trace amounts of heavy metals to the environment of microbial cells often stimulates microbial growth (Gikas, 2007; 2008).

However, higher concentrations result in severe reduction of microbial activity, which is reflected by reduction of apparent growth rate and increase in lag time. Heavy metal toxicity on microorganisms has been modelled by Utgikar *et al.* (2003) who studied the effects of Cu and Zn ions toxicity on sulphate reducing bacteria. The aim of this study was to assess the resistance and effect of different concentrations of Na+1, Ca+2 and Zn+2 on growth of *Bacillus* and *Pseudomonas* spp and to check effect of different incubation periods on resistance to metal by *Bacillus* and *Pseudomonas* spp.

MATERIALS AND METHODS

Sample collection

The bacterial samples were obtained from Research Laboratory, Department of Science Laboratory Technology,

Imo State Polytechnic Umuagwo, Nigeria. These bacterial samples were identified and categorized as heavy metal resistant isolates and then preserved on nutrient agar slants during previous work at GCUF (Umber, 2008).

Culturing of preserved heavy metal resistant bacteria

Fresh cultures of preserved bacterial isolates were obtained by growing them on nutrient agar.

Purification and preservation of heavy metal resistant bacteria

Colonies of *Bacillus* and *Pseudomonas* spp. were picked and grown again and again on nutrient agar plates to remove contaminations. These two genera (*Bacillus* and *Pseudomonas*) were separated on the basis of morphological and biochemical tests, and preserved on slants at about 8°C.

Biochemical Tests

Following biochemical tests were performed for the identification of bacteria. Gram's Staining, Catalase Test, Oxidase Test, Indole Test, Motility test, Vogas Proskauer Test.

Determination of effect of different concentration of metals (Zn+2, Na+1, Ca+2) on growth of bacteria Oladipo *et al.*, (2010).

100 ml of distilled water was taken in 250 ml flasks, and added 2.5 grams of nutrient broth in it. Metallic salts ZnCl_2 , NaCl, CaCl_2 were used as source of Metals ions. These Metals ions (Zn^{2+} , Na^{+1} , Ca^{2+}) were used in varying concentrations 25, 50, 75, 100, 125, 150, 175, 200 mg/ml, added these concentration in nutrient broth solution. The pH of the solutions was checked by pH meter and heats them till boiling. They were sterilized for 15 minutes at 121°C in autoclave. After cooling, the flasks were inoculated with the test organism under laminar flow and incubated at 37°C in incubator on shaker at 160 rpm. Growth was detected using atomic absorption spectrophotometer in terms of optical density at 600 nm after 0, 24, 48, 72, 96 hours of incubation time. Increase in turbidity of the medium was recorded as positive for growth while a negative result showed no turbidity. Flasks without metals served as control. All the experiments were conducted in triplicates.

Statistical Analysis

The data was subjected to statistical analysis in which uptake of various metals (Na^{+1} , Ca^{2+} , Zn^{2+}) by *Bacillus* and *Pseudomonas* spp. at pH 7 and temperature 37 °C was compared using student's t test.

RESULTS

Identification of Bacterial Isolates

On the basis of morphological and biochemical studies two bacterial genera were isolated namely *Bacillus* and *Pseudomonas* spp.

Bacillus sp. and Metal resistance pattern

Na+1: Figure 1 shows effect of Sodium metal on growth of *Bacillus* spp. cultivated at 37°C and pH 7. A significant increase in growth with time of incubation with metal

Table 1: Morphological and biochemical results

Bacteria identified	<i>Bacillus</i> spp.	<i>Pseudomonas</i> spp.
Colony	Creamy white	Creamy white
Morphology	irregular	round
Gram's staining	+Ve	-Ve
Catalase test	+Ve	+Ve
Oxidase test	+Ve	+Ve
Indole test	+Ve	-Ve
Motility test	+Ve	+Ve
Voges Proskauer test	+Ve	-Ve

Key: +ve = Positive; -ve= Negative

concentrations ranged between 25-75 mg/ml was observed. However, very little increase in growth was observed with Sodium metal concentration ranged between 100-150 mg/ml. The concentration of 175 mg/ml of Na+1 metal had no effect on *Bacillus* spp. growth. Whereas, a gradual decrease in growth of isolate was found at 200 mg/ml. Moreover, *Bacillus* sp. Showed increase in growth up to 72 hours of incubation with almost all Na+1 concentration except 200 mg/ml, while death phase occurred at 96 hours.

Ca₂+: It was noticed that 25, 50, 75 and 100 mg/ml of Ca₂+ had a positive effect on *Bacillus* sp. and increased the growth from 24 to 72 hours of incubation, afterwards at 96 hours, growth significantly decreased. While, 125, 150, 175 and 200 mg/ml of Ca₂+ had a negative effect and decreased the growth with increase in time of incubation (Fig. 2).

Zn₂+: It was observed that 25 and 50 mg/ml of Zn₂+ of metal concentration in the medium had a positive effect on *Bacillus* spp. It caused increase in the growth significantly till 72 hours of incubation while, decrease after 96 hours of incubation. Zn₂+ metal concentrations of 75 and 100 mg/ml in medium had no effect on *Bacillus* sp. However, at the concentration of 125, 150, 175 and 200 mg/ml of Zn₂+, growth decreased with increase in time of incubation (Fig. 3).

***Pseudomonas* sp. and Metal resistance pattern**

Na+1: A significant and constant raise in growth of *Pseudomonas* spp. was noticed at 25 and 50 mg/ml concentration of Na+1 metal even after 96 hours of incubation. In addition, growth of bacterial isolate increased at 75 and 100 mg/ml of Na+1 metal concentration up to 72 hours of incubation, afterwards it declined. However, no effect of Na+1 metal concentration of 125 and 150 mg/ml were observed. A further increase of metal concentration (175 and 200 mg/ml) had caused gradual decline in the growth of *Pseudomonas* spp. (Fig. 4).

Ca+2: It was noticed that 25 and 50 mg/ml of Ca₂+ had a positive effect on *Pseudomonas* spp. and increased the growth from 24 to 72 hours, afterwards growth significantly decreased at 96 hours of incubation. At 75-125 mg/ml concentration of Calcium metal, growth remained unaffected. Whereas, the concentrations of 150, 175 and 200 mg/ml of Ca₂+ Metal ion had a negative effect and decreased the growth.

Zn+2: It was observed that 25 and 50 mg/ml concentrations of Zn₂+ had a positive effect on




Fig. 1: Effect of different concentrations of Na+1 on growth of *Bacillus* sp. on different incubation periods.




Fig. 2: Effect of different concentrations of Ca₂+ on growth of *Bacillus* spp. on different incubation periods.




Fig. 3: Effect of different concentrations of Zn₂+ on growth of *Bacillus* spp. on different incubation periods.




Fig. 4: Effect of different concentrations of Na+1 on growth of *Pseudomonas* spp. on different incubation periods.




Fig. 5: Effect of different concentrations of Ca₂+ on growth of *Pseudomonas* spp. on different incubation periods.

Pseudomonas sp., it increased the growth significantly till 72 hours but decreased after 96 hours of incubation. At 75 mg/ml of Zn₂+ growths remained unaffected till 72 hours

afterwards it decreased. In addition, 100 - 200 mg/ml of Zn^{2+} decreased the growth with increase in time of incubation. It could be concluded that this *Pseudomonas* spp. can tolerate Zn^{2+} up to 75 mg/ml.

Comparison of effects of metals between *Bacillus* spp. and *Pseudomonas* spp.

Figures 7, 8 and 9 Show that *Bacillus* spp. was more resistant to Na^{+1} , Ca^{2+} and Zn^{2+} as compared to *Pseudomonas* spp.

Comparison of effect of heavy metal (Zn^{2+}) with two non heavy metals (Na^{+1} , Ca^{2+})

It was observed during the present investigation the minimum uptake of metal take place for Sodium metals as compared to other two metals; Ca^{2+} and Zn^{2+} . It was also found that highest growth values obtained in case of Calcium. This shows order of decreasing resistance which is as follows; Calcium > Zinc > Sodium.

DISCUSSION

In the present study 6 samples of preserved bacterial isolates were collected from research laboratory of Department of Science Laboratory Technology, Imo State Polytechnic Umuagwo, Nigeria. These bacterial isolates were originally isolated from industrial effluents of ALEX. In the present study, these bacterial isolates were re-identified and re-classified into two bacterial genera namely *Bacillus* and *Pseudomonas* spp. (Umber (2008). The present study was carried out to explore the effects of Na^{+1} , Ca^{2+} , Zn^{2+} , on growth of *Bacillus* and *Pseudomonas* spp. In the present study re-identified, purified and characterized isolates were preserved on fresh nutrient agar slants, stored at $4^{\circ}C$ for 24 hours and later incubated at $37^{\circ}C$ for 24 hours prior to each study. Nutrient broth was used for the cultivation of *Bacillus* and *Pseudomonas* spp. This was also reported by Ibezim *et al.* (2006). These authors discovered that *Bacillus* and *pseudomonas* spp. grew well between 24 to 72 hours of incubation period. Prescott *et al.* (2005) and Ynte *et al.* (2004) also agreed with the findings of Ibezim *et al.*, (2006). Increase in turbidity of the medium was recorded as positive for growth while a negative result shows no turbidity as reported by Oladipo *et al.*, (2010).

The use of NaCl salt as a source of Na^{+1} metal is also reported by Abdulkarim (2009) and Oladipo *et al.* (2010). Decrease in growth of *Bacillus* and *Pseudomonas* spp. with increase in Na^{+1} metal concentration found in present study, has also been recorded by Komives *et al.* (2005); Okanlawon *et al.* (2010); Oladipo *et al.* (2010); Veenagayathri and Vasudevan (2010). The decrease in total growth due to increase in the salt concentration was probably as a result of hyper osmotic effect.

During the present study Calcium Chloride supplementation was found to have a positive influence on *Bacillus* and *Pseudomonas* spp. in terms of growth rate at lower concentrations (25-100 mg/ml) as previously mentioned by Saranya and Shenbagarathi (2010) and Yuangklang *et al.* (2010). While higher concentrations of Calcium (125 to 200 mg/ml) had a negative effect on growth.




Fig. 6: Effect of different concentrations of Zn^{2+} on growth of *Pseudomonas* spp. on different incubation periods.




Fig. 7: Comparison of effects of Na^{+1} between *Bacillus* and *Pseudomonas* spp.




Fig. 8: Comparison of effects of Ca^{2+} between *Bacillus* and *Pseudomonas* spp.




Fig. 9: Comparison of effects of Zn^{2+} between *Bacillus* and *Pseudomonas* spp.




Fig. 10: Comparison of effect of heavy metal (Zn^{2+}) with two non heavy metals (Na^{+1} , Ca^{2+}) on *Bacillus* spp.




Fig. 11: Comparison of effect of heavy metal (Zn²⁺) with two non heavy metals (Na⁺, Ca²⁺) on *Pseudomonas* spp.

Zinc is an essential trace element for growth and enzyme activities of bacteria. However, excess load of Zn shows toxicity and inhibition to microbial processes. In Nigeria, several zinc resistant strains were isolated. *Pseudomonas aeruginosa* CMG 103 was isolated from a metal polluted river and displayed a high level of zinc and cadmium stress. Similarly, two Zinc resistant strains resistant to ZnSO₄ up to 12.5 mg/ml were isolated by Hasnain and Sabri (1992) from industrial effluents. These isolates also showed tolerance to other heavy metals such as Ni²⁺, Ba²⁺, Fe³⁺ and Cd²⁺. ZnCl₂ was used as a salt supplement during the present research work and it was found that *Bacillus* and *Pseudomonas* spp. were resistant to low concentrations of Zinc (25-100 mg/ml) and growth increased with increase in time of incubation. However, higher concentrations (125-200 mg/ml) decreased the growth of both bacterial isolates. Similar findings were reported by El-Bstawy (2000) and Bong *et al.* (2010).

Account on comparison of effects of Na⁺, Ca²⁺, Zn²⁺ on growth of *Bacillus* and *Pseudomonas* spp., it was found that calcium had a positive effect on growth while sodium had negative effect on growth. Similar results were reported by Tarakci *et al.* (2004); Gomaa and Azab (2007); Alshiyab *et al.* (2008). During the present study it was also found that *Bacillus* spp. was more resistant to metals under investigation (Na⁺, Ca²⁺, Zn²⁺) as compared to *Pseudomonas* spp., this was also reported by Leung *et al.* (2000); Ravikumar *et al.* (2007).

REFERENCES

- Abdulkarim SM, AB Fatimah and JG Anderson, 2005. Effect of salt concentrations on the growth of heat stressed and unstressed *Escherichia coli*. J Food, Agric and Environ, 7: 51-54.
- Alshiyab H, MS Kalil, AA Hamid and WMW Yusuff, 2008. Trace metal effect on Hydrogen production using *C. acetobutylicum*. Online J Biol Sci, 8: 1-9.
- Azzaoui S, El Hanbali M and M Leblance, 2002. Copper, Lead, Iron and Manganese in the Sebou drainage basin; sources and impact on surface water quality. Water Qual Res J Can, 37: 773-784.
- Barefoot, R Robert, J Carl and MD Reich, 2002. The Calcium Factor: The Scientific Secret of Health and Youth, (6th ed). Wickenburg, AZ: Deonna Enterprises Publications: 447-448.
- Bong CW, F Malafatti, F Azam, Y Obayashi and S Suzuki, 2010. The effect of Zinc exposure on the bacteria abundance and proteolysis Activity in seawater. Interdisciplinary Studies on Environmental Chemistry Biological Responses to Contaminants, Eds, 8: 57-63.
- Chindah AC, AS Braide and OC Sibeudu, 2004. Distribution of hydrocarbons and heavy metals in sediment and a crustacean (shrimps *Penaeus notialis*) from the bonny/new Calabar river estuary, Niger Delta. Afric J Environ Assess, Manage, 9: 1-14.
- El-Bstawy E, 2000. Selective bioaccumulation of mixed heavy metals by a range of brackish water bacteria. Pak J Biol Sci, 3: 1496-1499.
- Emongor V, E Kealotswe, I Koorapetse, S Sankwasa and S Keikanetswe, 2005. Pollution indicator in Gaberone effluent. J Appl Sci, 5: 147-150.
- Ensink JHJ, HW Vander, Y Mastsuno, S Munir and MR Aslam, 2002. The use of untreated waste water in Peri-urban Agriculture in Pakistan: risks and opportunities. Research Report No. 64, International Water Management Institute, Colombo, Sri Lanka, 22pp.
- Gikas P, 2007. Kinetic responses of activated sludge to individual and joint nickel (Ni (II)) and cobalt (Co (II)) - an isobolographic approach. J Hazard Mater, 14: 246-256.
- Gikas P, 2008. Single and combined effects of nickel (Ni(II)) and cobalt (CO(II)) ions on activated sludge and on other aerobic microorganisms. A review, J Hazard Mater, 159: 187-203.
- Gomaa OM and KH Azab, 2007. Role of Calcium Carbonate in protecting the colony forming ability of *Bacillus simplex* TWW-04 exposed to oxidative stress. J Adv Biol Res, 1: 49-55.
- Hasnain S and AN Sabri, 1992. Effects of temperature and pH on conjugal transfer of Zinc resistant plasmids residing in Gram negative bacteria isolated from industrial effluents. J Environ Pollut, 76: 245-249.
- Ibezim EC, CO Esimone, O Okorie, IV Onyishi, PO Nnamani, SA Brown and CE Obodo, 2006. A study of the in vitro interaction of Cotrimoxazole and Ampicillin using the checkerboard method. Afric J Biotech, 5: 1284-1288.
- Komives CF, LY Cheung, SB Pluschkell and MC Flickinger, 2005. Growth of *Bacillus methanolicus* in sea water based media. J Indust Microbio, 32: 61-66.
- Latif MI, RA Sial, MF Chaudhary, ST Abbas and AG Khan, 2008. Quality of effluents from Hattar Industrial Estate. J Zhejiang Uni Sci, 7: 974-980.
- Leung WC, MF Wong, W Lo, HF Yup and CK Leung, 2000. Removal and recovery of heavy metals by bacteria isolated from activated sludge treating industrial effluents and municipal waste water. J Biotech, 41: 233-240.
- Minna M, JG Jussila, K Lindstrom and L Suominen, 2006. Genetic diversity of culturable bacteria in oilcontaminated rhizosphere of *Galega orientali*. J Environ Pollut, 139: 244-257.
- Okanlawon BM, ST Ogunbanwo and AO Okunlola, 2010. Growth of *Bacillus cereus* isolated from some traditional condiments under different regimens. Afric J Biotech, 8: 2129-2135.
- Oladipo LC, DT Adeleke and AO Adebisi, 2010. The effect of pH and chemical preservatives on the growth of bacterial isolates from some Nigerian packaged fruit juices. Pak J Biol Sci, 13: 16-21.
- Pandey G and RK Jain, 2002. Bacterial chemotaxis towards environmental pollutants. J Appl Environ Microbio, 68: 5789-5795.

- Prescott LM, PJ Harley and AD Klein, 2005. Microbiology. (6th Edition). McGraw-Hill Companies, Inc, New York, USA: 951pp.
- Rani G, 2003. Microbial biomass: An economical alternative for removal of heavy metals from wastewater. Indian J Exp Biol, 41: 945-966.
- Ravikumar S, GP William, S Shanthi, NAA Garcelin, S Babu and PS Parimala, 2007. Effect of heavy metals (Hg and Zn) on the growth and Phosphate solubilising activity in Halophilic Phosphobacteria isolated from Manakudi mangrove. J Environ Biol, 28: 109-114.
- Sani RK, BM Peyton and M Jandhyala, 2003. Toxicity of Lead in aqueous medium to *Desulfovibrio desulfuricans* G20. J Toxicol Environ Chem, 22: 252-260.
- Saranya V and R Shenbagarathai, 2010. Effect of Nitrogen and Calcium sources on growth and production of PHA of *Pseudomonas* sp. LDC-5 and its mutant. Current Res Biol Sci, 2: 164-167.
- Tarakci Z, E Sagun, H Sancak and H Durmaz, 2004. The effect of salt concentration on some characteristics in herby cheese. Pak J Nutr, 3: 232-236.
- Tariq J, M Ashraf, M Jaffar and M Afzal, 2006. Pollution status of Indus River Pakistan through heavy metal and micronutrient content of fish, sediment and water. J Water Res, 30: 1337-1344.
- Ugochukwu CNC, 2004. Effluent monitoring of oil servicing company and its impact on the environment. Afric J Environ Assess, Manage, 8: 27-30.
- Umber L, 2008. Isolation of Heavy metal resistant bacteria from different industrial effluents of Faisalabad and Jhang. MSc Thesis. Department of Zoology, Government College University, Faisalabad, 34-98.
- Utgikar VP, HH Tabak, JR Haines and R Govind, 2003. Quantification of toxic and inhibitory impact of Copper and Zinc on mixed cultures of sulphate reducing bacteria. J Biotechnol Bioeng, 83: 306-312.
- Valdman E, L Erijman, FLP Pessoa and SGF Leite, 2001. Continuous biosorption of Cu and Zn by immobilized waste biomass *Sargassum* sp. Process Biochemistry, 36: 869-873.
- Veenagayathri K and N Vasudevan, 2010. Effect of pH, Nitrogen sources and salts on the degradation of phenol by the bacterial consortium under saline conditions. J Biotechnol Biochem, 6: 783-791.
- Vutukuru SS, 2003. Chromium induced alterations in some biochemical profiles of the Indian major carps, *Labeo rohita* (Hamilton). Bulletin Environ Contamination Toxicol, 70: 118-123.
- Ynte PDE, LM Vries, WM Hornstra and TADE Vos, 2004. Growth and population of *Bacillus cereus* ATCC 14579 under defined conditions: Temporal Expression of Genes for Key Sigma Factors. J Appl Environ Microbio, 70: 2514-2519.
- Yuanklang C, C Wachirapakorn, HE Mohamed, A Alhaidary and AC Beynen, 2010. Effect of Calcium supplementation on growth, nutrient digestibility and fecal *Lactobacilli* in dairy calves. J Anim Vet Sci, 5: 127-131.