



Research Article

Initial Seed Moisture Content, Storage Temperature and Storage Duration Influenced Arabica Coffee (*Coffea arabica* L.) Seedling Performances

Kalifa Nasiro

Ethiopian Institute of Agricultural Research, Jimma Agricultural Research Center, P.O. Box 192, Jimma, Ethiopia

*Corresponding author: nasirokalifa@gmail.com

Article History: 888 Received: September 16, 2020 Revised: November 16, 2020 Accepted: December 18, 2020

ABSTRACT

Coffee is one of the most important agricultural products in the international market and many countries are involved in its production, trade or consumption. Arabica coffee is self-pollinated and homozygous, they are normally propagated by seeds. Seeds have been considered intermediate storage behavior with varying results. It is highly desirable that seeds are stored safely to optimize coffee seedling production at the appropriate time and season with ideal climatic conditions for planting in the field. The objective of this study was to evaluate the effect of storage temperature, time of storage and initial seed moisture content on early seedling performances of coffee seeds and to determine the appropriate seed handling method. In this experiment, the influence of initial seed moisture content with four levels (12, 17, 22 & 27%) storage temperature (ST) with two levels (15°C & ambient), time of storage with six levels (sowing after 1,2,3,4,5 & 6 months) and on coffee seedling growth stages were studied in a split-split-plot factorial design. The data collected were subjected to Analysis of Variance (ANOVA) using statistical analysis system version 9.2 software (SAS, 2009). Treatment means were separated using LSD at 0.05 probability level. The present findings of storage environment with cold temperature (15°C) accelerated seedling growth much better than did ambient temperature condition. All tested seedling growth stages were faster at initial time of storage. After third month of storage, seed quality drastically reduced especially under ambient storage condition. Seeds dried to 12% moisture content showed delay of germination and growth throughout the trial period. Seeds with 27% initial moisture content took shorter time to reach at different growth stages at initial storage time but when aged took much time. Storage temperature, time of storage and initial seed moisture contents showed highly significant main and interaction effects and seeds dried to intermediate moisture level (17 & 22%), stored under cold temperature and sown at early times resulted in enhanced seedling growth. Hence, the current study suggests drying coffee seeds to 17% to 22% moisture contents and store under relatively lower temperatures at about 15°C for not more than six months of storage. As the present finding was limited to single cultivar and specific environmental condition further investigation is significant.

Key words: Arabica coffee, Seed Storage, Seedling growth stages.

INTRODUCTION

The Coffee seed normally presents high germination potential, just after appropriate harvest and desiccation. However, it loses its physiological performance very rapidly under usual storage conditions. Ellis *et al.* (1990) have grouped coffee under intermediate category for seed behavior, because seed viability lasts for only short periods and must be planted soon after drying. But one may need to store seeds for up to six months or more because of the gap between harvesting (during October and November) and sowing seeds (between April and

August) for Ethiopian seeding calendar. Hence, storage and maintenance of quality seed lots until sowing is as important as producing quality seeds.

Jimma agricultural research center (JARC), which is the sole mandated coffee seed producer in the country, is conventionally drying to 12 to 15% seed moisture ranges for seed storage. However, feedback from various users indicating that there is a problem on germination. This justifies further investigation on appropriate handling of coffee seed in order to come-up with the best solution for above-mentioned problems. Therefore, due to the limited information in our country about the storability of the seeds

Cite This Article as: NasiroK, 2020. Influence of seed initial moisture Content, Storage Condition and time of storage on seedling growth stages of coffee (*Coffea arabica* L.). Int J Agri Biosci, 9(6): 278-283. www.ijagbio.com (©2020 IJAB. All rights reserved)

the current study is justified with the aim of giving practical information about handling and storing of coffee seeds.

Objectives

To evaluate the effect of initial seed moisture content, storage temperature, and duration of storage on coffee seedling growth performances and to determine the appropriate seed handling method.

MATERIALS AND METHODS

Description of the study site

The experiment was executed in a nursery site at Jimma Agricultural Research Center (JARC), of the Ethiopian Institute of Agricultural Research. It is the National Coffee Research Coordinating Center in the country and is located 365 km away from Addis Ababa, 12 km from the Jimma town in the south west direction. The center is found within tepid to cool humid highland agro-ecological zone of the country at an altitude of 1750 meters above sea level, 7° 46' N, latitude, and 36°47' E longitude in the sub humid tropical belt of south western Ethiopia. Average annual rain fall of the area is 1594 mm with 67% mean relative humidity. The mean minimum and maximum temperatures are about 11.6°C and 26.3°C respectively. The soil of the center has a characteristic of reddish to reddish brown clay nature, where eruticnitosols and chronic cambisols are dominant types with pH range of 5 to 6.

Experimental materials

Following the conventional procedures widely applied in coffee seed preparation (Van der Vossen, 1979; Goodman, 1980; Rothofs, 1980), fully ripe red cherries were harvested from selected mother trees in the seed orchard of cultivar 74110 (which was selected as it is widely adapted, highly demanded and much produced) at JARC, on 21st November 2011 growing season. This cultivar was selected for the present study, because it is high-yielding CBD resistant, widely adaptable, highly demanded and produced much. The cherries were sorted out and pulped in a hand pulp separator. After pulping the selected cherries, the wet parchment beans were again sorted out, thoroughly washed, and taken to drying room, hut made of grass roof.

Then the wet parchment coffee was laid on wire mesh for drying under shade and when its skin dried dressed in fine wood ash following the JARC conventional practice and kept till it attained the desired four levels of moisture contents (27, 22, 17 and 12% [fresh weight base]). One kg of parchment coffee seeds was taken on 5th January 2012 till 2nd July 2012 from each of the four batches and kept separately in each of the two storage conditions. The storage conditions used in this study were cold store with 15°C (SC1) and a room at ambient temperature (SC2). All the seed lots with four levels of MCs were kept under both storage conditions.

Representative samples of seeds were taken every month from each treatment combinations and were subjected to a series of tests in the laboratory and nursery trials to evaluate the potential viability (germination) and early seedling growth potential status of each seed lot.

Treatments and experimental design

A split-split plot factorial design was used with three replications.

Factors

- A. Storage condition (cold 15°C & ambient temperature)
- B. Time of storage (sowing after 1, 2, 3, 4, 5 & 6 months of storage)
- C. Initial seed moisture content (12%, 17%, 22% & 27%)

As presented in Table1, in this experiment, storage conditions (SC) was assigned to main plot, time of storage (ST) was assigned to sub-plots while seed initial moisture level was assigned to sub-sub-plots. Total number of treatment combinations were 48 (2*6*4) replicated three times and the total number of experimental plots were 144. The treatments were randomly and independently assigned to main plots, sub plots and the sub-sub plots. Every routine nursery activity was practiced uniformly to all experimental units as per the recommendation of the JARC (Institute of Agricultural Research, 1996).

The model

Three factor analysis of variance model was used with General Linear Model (GLM) Procedures of SAS Version 9.2. The linear statistical model for the split-split-plot design would be:

$$Y_{ijk} = \mu + \alpha_i + \beta_j + \gamma_k + (\alpha\beta)_{ij} + (\alpha\gamma)_{ik} + (\beta\gamma)_{jk} + (\alpha\beta\gamma)_{ijk} + \epsilon_{\mu_{ijk}}$$

Where: -

Y_{ijk} = the response measurement for the ijk th observation
 μ = is the overall mean effect.

α_i = is the effect of the i th level of sub-sub-plot

β_j = is the effect of the j th of main plot

γ_k = is the effect of the k th of sub-plot

$(\alpha\beta)_{ij}$ = is the effect of the interaction b/n actor A&B

$(\alpha\gamma)_{ik}$ = is the effect of the interaction b/n factor A&C

$(\beta\gamma)_{jk}$ = is the effect of the interaction b/n factor C&B

$(\alpha\beta\gamma)_{ijk}$ = the effect of interaction b/n the three factors

$\epsilon_{\mu_{ijk}}$ = is a random error component for all factors

Sampling and data collection

The observations were recorded on seed germination percent, shoot length, seedling root length, girth diameter, leaf area, seedling dry weight and seedling vigor Index and they were explained below.

Determination of percentage of seedling growth stages

The plots were checked consistently every five days starting from the 65th day after sowing till all the potentially capable seedlings in each plot developed their cotyledonary leaves, total percentage of seedlings at the butterfly growth stage (%BFS) was determined for each plot. Using the number of seedlings that produced their cotyledonary leaves, or that attained the "butterfly" stage of growth, the mean days required to attain this stage of growth (MDBFS) was also calculated following the same procedure used for determination of MDE above.

Similarly, the number of seedlings that developed their pair of true leaves were further counted from each plot and recorded at five-days interval starting from the 80th day after sowing and the percentage of seedlings that

produced their first true leaf (%TLS) was determined. The mean number of days required by each plot to attain this growth stage (MDTLS) was also computed using the method described above.

Statistical analysis

The seedling growth data collected were subjected to Analysis of variance (ANOVA) for split-split-plot design and treatment mean separation was carried out using least significant difference (LSD) at 0.01 and 0.05 probability levels using General Linear Model (GLM) procedure of SAS statistical version 9.2 software (SAS, 2009).

RESULTS AND DISCUSSION

Butterfly Growth Stage (BFS %)

The data of the experiment showed that the number of seedlings attained butterfly growth stage was significantly ($P < 0.01$) influenced by the main and interaction effects of storage condition, storage time and seed moisture content (Table 1). Treatment combinations of seeds stored in cold condition dried to 22% moisture contents and sown at first month maintained significantly higher butterfly growth stage (88.00%). Similarly, seeds stored in cold condition dried to 12%, 17% & 27% moisture contents and sown at first month and seeds with 17%, 22% & 27% moisture contents that sown at second month also maintained higher percentage of butterfly growth stage (Table 2). This indicated that cold condition better maintained coffee seeds quality and it is consistent

with the findings of Barboza and Herrera, (1990); Ellis *et al.*, (1990, and 1991) who have proposed the use of low temperatures (around 15°C) to preserve coffee seed viability and vigor for a considerable period of time.

The least butterfly growth stage was recorded in the seeds stored in ambient condition dried to 12% & 17% moisture contents and sown at sixth month of storage (31.00 & 32.67%, respectively) (Table 2). This could be attributed to the general decline in viability of seed lots as a result of prolonged storage. Beside failure in germination, which is a sign of seed death, the general symptoms of deterioration exhibited by seeds include: reduced germination rate and vigor, and development of abnormal seedlings (Decouche and Caldwell, 1960; Anderson, 1970; Negussu, 1986).

Initial seed moisture contents with 22% maintained better performance for percent butterfly stage with prolonged storage time. These results are in accordance with the reports of (IAR, 1996; and Wondyifraw, 1994).

True Leaf Growth Stage (TLS %)

Interaction effects of storage condition, storage time and seed moisture content on true leaf growth stage was significant ($P < 0.01$) similarly with the results of emergence percentage butterfly stage (Table 1). Treatment combinations of seeds stored in cold condition dried to 17% moisture content and sown after a month maintained significantly higher true leaf growth stage of (85.67%). However, not significantly differed from seeds stored under cold condition that dried to 22 and 27% seed

Table 1: Analysis of Variance for butterfly stage (%BFS), true leaf stage (%TLS), mean days to butterfly (MDBFS) and true leaf stage (MDTLS)

Source of Variation	DF	BFS	TLS	MDBFS	MDTLS
Rep	2	73.42**	85.33**	5.76**	6.30*
SC	1	9136.17**	10885.44**	680.34**	1778.03**
Error (a)	2	26.76**	21.36*	11.17**	22.13**
ST	6	3732.66**	4021.23**	523.62**	1390.04**
SC*ST	6	870.99**	830.33**	24.72**	79.06**
IMC	3	179.58**	156.75**	26.43**	54.43**
SC*IMC	3	15.90*	30.35**	5.78**	10.94**
ST *IMC	18	14.23**	14.44**	3.64**	2.09 ^{ns}
SC*ST*IMC	18	13.33**	17.59**	0.90*	6.06**
Error (b)	12	22.68**	25.00**	7.62**	9.13**
CV%		2.89	3.51	0.81	0.89

*, ** = Indicate significant differences at the 5% and 1% probability levels, respectively. ns = non-significant at 5% probability levels.

Table 2: The interaction effects of storage condition (SC), storage time (ST) and initial seed moisture content (IMC) on percentage of seedlings attained butterfly stage (%)

SC	IMC	Storage Time (in month)					
		1	2	3	4	5	6
Cold (15°C)	12%	83.67 ^{a-e}	80.33 ^{d-i}	77.33 ^{g-l}	72.33 ^{m-p}	71.33 ^{nop}	68.67 ^{pq}
	17%	86.67 ^{ab}	83.33 ^{a-f}	80.00 ^{d-i}	75.00 ^{j-n}	72.33 ^{m-p}	66.33 ^q
	22%	88.00 ^a	84.33 ^{a-d}	82.00 ^{b-g}	80.33 ^{d-i}	76.67 ^{i-m}	71.33 ^{nop}
	27%	87.67 ^a	86.00 ^{abc}	81.00 ^{d-i}	77.33 ^{g-l}	73.00 ^{l-p}	69.00 ^{pq}
Ambient	12%	81.67 ^{c-h}	78.00 ^{g-k}	72.67 ^{l-p}	55.33 ^f	41.33 ^u	31.00 ^w
	17%	80.67 ^{d-i}	78.67 ^{f-k}	74.00 ^{k-o}	54.33 ^{rs}	49.00 ^t	32.67 ^{vw}
	22%	82.00 ^{b-g}	79.33 ^{e-j}	77.00 ^{h-m}	65.33 ^q	49.67 st	41.67 ^u
	27%	82.00 ^{b-g}	74.67 ^{j-o}	70.00 ^{opq}	56.33 ^r	47.00 ^t	37.33 ^{uv}
CV%		4.12					
LSD (5%)		4.6806					

Means followed by the same letter(s) are not significantly different at 5% level of probability.

moisture content and sown after a month and with the same three levels of moisture contents that sown after second month. While, the least percentage of true leaf growth stage was recorded in the seeds stored in ambient condition dried to 12 and 17% moisture contents and sown after sixth month of storage (27.67%). Similarly, seeds stored in ambient condition dried to 22 and 27% moisture contents and sown after sixth month of storage also showed lower percentage of seedlings attained true leaf stage (Table 3). This could be attributed to the general decline in viability of seed lots as a result of prolonged storage.

In the same manner to that of butterfly stage seeds with 22% moisture content maintained better performance for percent true leaf stage with prolonged storage time. These results are in accordance with the reports of (IAR, 1996; and Wondyifraw, 1994). The general symptoms of deterioration exhibited by seeds include: reduced germination rate and vigor, and development of abnormal seedlings (Decouche and Caldwell, 1960; Anderson, 1970; Negussu, 1986).

Mean Days to Butterfly Stage (MDBFS)

The data of the experiment showed interaction between storage condition, time of storage and initial seed moisture content was highly significant ($P < 0.01$) for mean days to butterfly stage (Table 1). The treatment combinations of seeds dried to 27%, stored under cold

condition and sown after a month showed significantly short MDBFS (87.67 days). While, extended time was recorded for treatment combination of seeds stored in ambient condition, dried to 12% moisture content and sown after the end of the sixth month of storage (107 days). This could be attributed to the general decline in viability and vigor of seed lots as a result of prolonged storage, fluctuation of temperature and high respiration under ambient condition. The result of this experiment revealed that seeds dried to 22% moisture and stored in cold condition better maintained rapidity and uniformity of seedlings than did the other treatment combinations (Table 4). These results agree with the reports of (Van der Vossen, 1979; Bradford *et al.*, 1990; Ellis *et al.*, 1990; and Rosa, *et al.*, 2005) who worked on various crops including coffee.

Mean days to butterfly stage progressively declined with prolonged storage time irrespective of moisture content, storage environment and their interaction. These results are in accordance with the reports of (IAR, 1996; and Wondyifraw, 1994). There was a positive relationship between the mean number of days required to reach a given growth stage and time in storage. Vigorous seeds rapidly germinate, metabolize and establish in the field. Therefore, any method used to determine the rapidity of growth of the seedling will give an indication of seed vigour level (Perry, 1984).

Table 3: The interaction effects of storage condition (SC), storage time (ST) and initial seed moisture content (IMC) on percentage of seedlings attained at true leaf stage (%)

SC	IMC	Storage Time (in month)					
		1	2	3	4	5	6
Cold (15°C)	12%	79.33 ^{b-e}	75.00 ^{e-j}	73.67 ^{f-l}	69.67 ^{k-o}	69.33 ^{k-p}	64.33 ^{p-s}
	17%	85.67 ^a	80.67 ^{a-d}	76.67 ^{c-g}	70.00 ^{j-o}	65.67 ^{o-r}	61.00 ^{r-s}
	22%	85.33 ^a	81.33 ^{abc}	78.67 ^{b-f}	75.00 ^{e-j}	71.00 ⁱ⁻ⁿ	67.67 ^{m-q}
	27%	85.00 ^a	83.33 ^{ab}	79.00 ^{b-e}	73.67 ^{f-l}	68.67 ^{l-q}	66.00 ^{n-r}
Ambient	12%	78.00 ^{c-f}	71.67 ^{g-m}	64.00 ^{q-s}	48.67 ^{tu}	36.33 ^w	27.67 ^x
	17%	76.67 ^{c-g}	75.67 ^{d-i}	71.33 ^{h-m}	51.00 ^t	43.67 ^{uv}	27.67 ^x
	22%	75.67 ^{d-i}	76.33 ^{c-h}	71.00 ⁱ⁻ⁿ	60.00 ^s	44.00 ^{uv}	32.67 ^{wx}
	27%	74.33 ^{e-k}	71.00 ⁱ⁻ⁿ	65.00 ^{o-s}	52.67 ^t	43.00 ^v	30.33 ^x
CV%	4.76						
LSD (5%)	5.0679						

Means followed by the same letter(s) are not significantly different at 5% level of probability.

Table 4: The interaction effects of storage condition (SC), storage time (ST) and initial seed moisture content (IMC) on mean days taken to seedlings to attain at butterfly stage (no of days)

SC	IMC	Storage Time (in month)					
		1	2	3	4	5	6
Cold (15°C)	12%	90.00 ^{bcd}	91.00 ^{b-f}	93.00 ^{f-i}	95.67 ^{j-m}	98.33 ^{n-q}	101.33 ^{stu}
	17%	91.33 ^{c-f}	93.00 ^{f-i}	94.33 ^{h-k}	95.00 ^{i-l}	97.67 ^{m-p}	99.33 ^{p-s}
	22%	89.00 ^{ab}	91.00 ^{b-f}	92.00 ^{d-g}	93.67 ^{g-j}	95.67 ^{j-m}	97.67 ^{m-p}
	27%	87.67 ^a	90.67 ^{b-e}	92.67 ^{e-h}	94.67 ^{h-k}	96.00 ^{klm}	100.00 ^{q-t}
Ambient	12%	93.00 ^{f-i}	95.00 ^{i-l}	97.67 ^{m-p}	101.00 ^{r-u}	104.33 ^{wx}	107.00 ^y
	17%	92.00 ^{d-g}	95.00 ^{i-l}	96.33 ^{k-n}	99.00 ^{o-r}	102.67 ^{uvw}	105.33 ^{xy}
	22%	90.00 ^{bcd}	93.00 ^{f-i}	95.33 ^{jkl}	99.33 ^{p-s}	102.00 ^{tuv}	104.67 ^{wx}
	27%	89.67 ^{abc}	93.00 ^{f-i}	97.00 ^{l-o}	101.33 ^{stu}	104.00 ^{vwx}	107.33 ^y
CV%	1.40						
LSD (5%)	2.1862						

Means followed by the same letter(s) are not significantly different at 5% level of probability.

Table 5: The interaction effects of storage condition (SC), storage time (ST) and initial seed moisture content (IMC) on mean days taken to seedlings to attain at true leaf stage (no of days)

SC	IMC	Storage Time (in month)					
		1	2	3	4	5	6
Cold (15°C)	12%	119.67 ^{b-e}	120.67 ^{b-e}	123.67 ^{f-h}	127.67 ^{jmn}	131.67 ^{op}	135.00 ^{qr}
	17%	119.00 ^{a-d}	120.00 ^{b-e}	122.00 ^{e-h}	124.67 ^{h-k}	128.00 ^{lmn}	135.00 ^{qr}
	22%	118.67 ^{a-d}	120.00 ^{b-e}	122.33 ^{e-h}	125.33 ^{i-l}	130.00 ^{no}	136.33 ^{rs}
	27%	116.33 ^a	118.33 ^{abc}	120.67 ^{b-e}	124.00 ^{g-j}	126.67 ^{j-m}	129.33 ^{mno}
Ambient	12%	122.33 ^{e-h}	126.00 ^{i-l}	130.33 ^{no}	135.33 ^{qr}	140.33 ^{tuw}	145.33 ^w
	17%	121.00 ^{c-f}	123.67 ^{f-i}	127.67 ^{lmn}	133.67 ^{pqr}	138.67 st	143.00 ^{vw}
	22%	120.00 ^{b-e}	124.33 ^{h-k}	128.00 ^{lmn}	134.00 ^{pqr}	140.00 ^{tu}	143.00 ^{vw}
	27%	118.00 ^{ab}	121.33 ^{d-g}	127.00 ^{klm}	133.33 ^{pq}	142.00 ^{uv}	145.33 ^w
CV%		1.31					
LSD (5%)		2.7214					

Means followed by the same letter(s) are not significantly different at 5% level of probability.

Mean Days to True Leaf Stage (MDTLS)

The result the present study showed that interaction between storage condition, time of storage and initial seed moisture content was highly significant ($P < 0.01$) for mean days to true leaf stage with the same manner to MDBFS (Table 1). The treatment combinations of seeds dried to 27% moisture content, stored under cold condition and sown after a month showed significantly short MDTLS (116.33 days). While, extended time was recorded for treatment combination of seeds stored under ambient condition, dried to 12% & 27% moisture and sown at the end of the sixth month of storage (145.33 days). This could be attributed to the general decline in viability and vigor of seed lots as a result of prolonged storage, fluctuation of temperature and high respiration under ambient condition. The result of this experiment revealed that seeds dried to 27% moisture and stored in cold condition better maintained rapidity and uniformity of seedlings than did the other treatment combinations (Table 5). This evidenced that the advantage of having optimum initial seed moisture content in maintaining seed viability and vigor for a longer time (Agrawal, 1992). Moreover, Bradford *et al.* (1990) also stressed that such rapid early growth of seedlings were associated with high vigor, a measure of the inherent potential of seed lots to produce strong plants, which in turn is affected by moisture content of the seed to be sown.

Mean days to true leaf stage progressively declined with prolonged storage time irrespective of moisture content, storage environment and their interaction. These results are in accordance with the reports of (Robinson, 1964; IAR, 1973; and Wondyifraw, 1994). This could be attributed to the general decline in viability of seed lots as a result of prolonged storage. As reported by Perry, (1984) vigorous seeds rapidly germinate, metabolize and establish in the field and therefore, any method used to determine the rapidity of growth of the seedling will give an indication of seed vigour level.

Summary and conclusions

Coffee has enormous economic, social and environmental significance in Ethiopia. But despite its enormous genetic variability, favorable ecology for production and its importance in the national economy of the country, productivity of the crop remained much lower than released varieties yielding potential which primarily

associated to use of poor seed that result in poor seedlings causing poor establishment in the field.

The present study was carried out with the objective of determining the quality of early seedlings growth potentials of coffee produced from seeds stored for six months with 27, 22, 17 and 12% initial seed moisture contents under storage conditions at 15°C and ambient temperature at JARC using split-split-plot factorial Design.

According to the results obtained in this investigation, it was observed that early seedling growth parameters tested resulted to be statistically influenced by the main and interaction effects of storage conditions, storage time and initial seed moisture contents. After four more months of storage all the factors resulted in highly significant ($P \leq 0.01$) difference for almost all the variables tested. The present findings of storage environment with cold condition (15°C) maintained seed germination and early seedling growth parameters much better performances than did ambient condition. All tested seed and seedling growth parameters were highest at initial time of storage and declined progressively with storage time. After third month seed quality drastically reduced especially under ambient storage condition. Seeds dried to 12% moisture content showed inferior performance throughout the trial period.

Seed stored under cold condition recorded minimum qualitative losses with better seed quality parameters throughout the storage period as compared to seeds stored under ambient condition. Seeds stored in cold condition maintained viability and vigor for longer period with minimum deterioration in seed quality.

Seeds with 27% initial moisture content showed higher performances at initial storage time but when aged drastically declined. Seeds with 22 and 17% initial moisture contents had maintained better seed germination and early seedling growth performances throughout the study period. Seeds dried to 22% followed by 17% initial moisture contents maintained higher germination, seedling emergence and early seedling growth parameters and stored at relatively lower temperatures. If stored under ambient condition should not be stored for longer than two to three months.

Storage condition, time of storage and initial seed moisture contents showed highly significant interaction effects and seeds dried to intermediate moisture level (17 & 22%), stored under cold condition and sown at early

times resulted in enhanced early seedling growth and uniformity in seedling. Hence, for storing coffee seeds, it is advisable drying the seeds to about 17% to 22% moisture contents and keep under storage with relatively lower temperatures (at about 15°C) for not more than five or six months of storage is advisable.

Recommendations and Future line of work

For immediate sowing of coffee seeds higher seed moisture content of 27% may be used for better seed germination and early growth potentials that result in vigorous seedling.

But if it is to be used after storage for more than two months initial seed moisture content needs to be reduced to around 17% for maintaining better seed quality for about six months.

For prolonged coffee seed storage up to six months storage temperatures nearly 15°C advised to be used and seeds kept under ambient condition may be sown immediately or within two to three months otherwise preferable not to be used as seed material.

The experiments conducted so far in this area are not sufficient to draw a reliable conclusion. Since, the present study was done for a single cultivar (74110) and under Jimma conditions further experiment needs to be conducted for varieties at different environmental conditions with narrower ranges of seed initial moisture content and storage temperatures.

REFERENCES

- Agrawal PK, 1992. Seed storage and packaging. In: *Techniques in Seed Science and Technology*. (2nd edn.) Ed. by P.K. Agrawal and M. Dadlani. South Asian Publishers Pvt Ltd, New Dehli. pp. 159-169.
- Alemayehu Teshome, Esayas Kebede and Kassu Kebede, 2008. Coffee Development and Marketing Improvement Plan. In: *Proceedings of A National Work Shop Four Decades of Coffee Research and Development in Ethiopia*. 14-17 August 2007, EIAR, Addis Ababa, Ethiopia. pp. 375-381.
- Anderson JD, 1970. Metabolic Changes in partially dormant wheat seed during storage. *Plant Physiology*. 46: 605-608.
- Barboza, R. and J. Herrera. 1990. Seed vigor in coffee seeds and its relations to drying temperature, moisture content and storage conditions. *Agronomy*, 14: 1-7.
- Bradford KJ, Steiner JJ and SE Travatha, 1990. Seed priming influence of germination and emergence of pepper seed lots. *Crop Sci*. 30 (3): 718-721.
- Central Statistics Authority (CSA), 2001. Annual report on Ethiopian Economy. 1999/2000, CSA, Addis Ababa, Ethiopia.
- Deloche JC and WP Caldwell, 1960. Seed vigor and vigor tests. *Proc. Assoc. Seed Anal.* 50: 124-129.
- Ellis RH, TD Hong and EH Roberts, 1990. An intermediate category of seed storage behaviour I. *Coffee. J. Exp. Bot.* 41: 1167-1174.
- Ellis RH, TD Hong and EH Roberts, 1991. An intermediate category of seed storage behavior. II. Effects of provenance, immaturity, and imbibition on desiccation-tolerance in coffee. *J. Exp. Bot.* 42: 653-657.
- FAO/WFP, 2008. Special report FAO/WFP crop and food supply assessment mission to Ethiopia, 24 January 2008.
- Goodman JH, 1980. Coffee nursery management in Guidelines on coffee management. Unpublished manual. Extension, CIP/MCTD. pp. 1-16.
- Institute of Agricultural Research (IAR), 1996. Recommended production technologies for coffee and associated crops. Research extension task force of Jima Research Centre. Addis Ababa, Ethiopia.
- Mayne, R., Tola, A. and Kebede, G. 2002. Crisis in the birth place of coffee, Oxfam International research paper, Oxfam International.
- Negussu Legesse, 1986. Effect of climate, insulation and ventilation on effective temperatures in open seed stores in the tropics. MSc. Thesis. Univ. of Edinburgh, Scotland.
- Perry DA, 1984. A vigor test for seed of barley (*Hordeum vulgare*) based on measurement of plumule growth. *Seed Sci. and Techn.* 5: 709-719.
- Rosa SD, VF Brandão, DS Junior, EVR Von Pinho, AD Veiga. and LHC Silva, 2005. Effects of different drying rates on the physiological quality of Coffea canephora.
- Rothofs B, 1980. Coffee production. Hamberg: Gordian-Max-Rieck.
- SAS (Statistical Analysis System), 2009. SAS user's guide: statistics, 5th ed, Statistical Analysis System Software. Version 9.2. SAS Inst., Cary, NC.
- Van der Vossen HAM, 1979. Methods of preserving the viability of coffee seed in storage. *Seed Sci. Technol.* 7: 65-74.
- Wondyifraw Tefera, 1994. The influence of duration of storage, initial moisture content and type of container on the viability of coffee (*Coffea Arabica L.*) seeds. An MSc. Thesis, Alemaya University, Ethiopia.